

2923-105  
MC:jdh



#10

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of )  
Stefan GRIMM, et al. )  
Serial No. 09/890,202 ) Examiner: Quang Nguyen  
Filed: November 5, 2001 ) Group Art Unit: 1636  
For: METHOD FOR PRODUCING )  
ENDOTOXIN-FREE NUCLEIC )  
ACIDS AND THE USE THEREOF)

SUBMISSION OF TRANSLATED COPY OF PRIORITY APPLICATION


Assistant Commissioner for Patents  
Washington, D.C. 20231

Dear Sir:

Submitted herewith is a certified translated copy of German Patent Application No. DE 19903507.5, filed January 29, 1999, from which priority has been claimed in the above-referenced patent application.

Respectfully submitted,

By

  
Martha Cassidy  
Attorney for Applicants  
Registration No. 44,066  
ROTHWELL, FIGG, ERNST & MANBECK, p.c.  
Suite 800, 1425 K Street, N.W.  
Washington, D.C. 20005  
Telephone: (202) 783-6040

Enclosure(s):  
Declaration by Translator  
Priority Certificate  
Translated Application



UNITED STATES PATENT AND TRADEMARK OFFICE

I, Susan ANTHONY BA, ACIS,

Director of RWS Group plc, of Europa House, Marsham Way, Gerrards Cross, Buckinghamshire, England declare;

1. That I am a citizen of the United Kingdom of Great Britain and Northern Ireland.
2. That the translator-responsible for the attached translation is well acquainted with the German and English languages.
3. That the attached is, to the best of RWS Group plc knowledge and belief, a true translation into the English language of the accompanying copy of the specification filed with the application for a patent in Germany on 29 January 1999 under the number 199 03 507.5 and the official certificate attached hereto.
4. That I believe that all statements made herein of my own knowledge are true and that all statements made on information and belief are true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the patent application in the United States of America or any patent issuing thereon.

For and on behalf of RWS Group plc

The 11th day of April 2003



**FEDERAL REPUBLIC OF GERMANY**

**[Eagle crest]**

**Priority Certificate  
for the filing of a Patent Application**

**File Reference:** 199 03 507.5

**Filing date:** 29 January 1999

**Applicant/Proprietor:** Xantos Biomedicine AG, Planegg/DE

First applicant:

ROCHE DIAGNOSTICS GmbH, Mannheim/DE

**Title:** Method for preparing endotoxin-free nucleic acids or nucleic acids  
with reduced endotoxin content and the use thereof

**IPC:** C 07 H, G 01 N

**The attached documents are a correct and accurate reproduction of the original  
submission for this Patent Application.**

Munich, 26 March 2003

**German Patent and Trademark Office**

**The President**

[Seal of the German Patent  
and Trademark Office]

pp

[signature]

Hiebinger

**Method for preparing ndotoxin-fre nucleic acids or  
nucl ic acids with reduced endotoxin content and the  
use thereof**

---

5 The invention relates to a method for isolating and  
purifying nucleic acids and/or oligonucleotides from a  
biological sample, to the use of the isolated or  
purified nucleic acid and/or oligonucleotide for  
transfecting cells and also for the production of an  
agent for the treatment of genetic disorders, to a  
10 composition suitable for the isolation or purification  
method and also to the use of potassium acetate and a  
silica gel-like support material for isolating  
endotoxin-free nucleic acids and/or oligonucleotides or  
nucleic acids and/or oligonucleotides with reduced  
15 endotoxin content.

The quality of isolated nucleic acids is becoming  
increasingly important. Highly pure nucleic acid  
fractions, i.e. fractions from which, if possible, all  
20 other cell components such as, for example, endotoxins,  
have been removed, play a central part in gene therapy  
or in transfecting cells of eukaryotic or also  
prokaryotic origin. Consequently, in the past few years  
methods or measures which allow the isolation of  
25 nucleic acids from biological sample material with high  
purity have increasingly been published. The  
established methods essentially make use of the use  
[sic] of affinity and/or anion [lacuna] chromatography  
materials and also of non-ionic detergents or also  
30 diluted solutions of higher alcohols. For example,  
according to W095/21177 the fractions of interest are  
subjected to an affinity chromatography or a  
chromatography on an inorganic solid phase, the latter  
preferably in the presence of a non-ionic detergent, in  
35 order to remove endotoxins and are then further  
purified by means of anion exchange chromatography. A  
two-stage chromatography method of this kind, however,

is time- and material-consuming and therefore is more academically valuable. According to another method (WO95/21178) a complicated anion exchange chromatography is likewise absolutely necessary in order to remove residues of a complex salt solution added beforehand.

Furthermore, it has been known for some time that DNA plasmids from complex biological samples of eukaryotic or prokaryotic origin can be isolated by binding to silica gel in the presence of chaotropic salts such as, for example, guanidine hydrochloride (M.A. Marko et al., *Analyt. Biochem.* 121, (1982) 382-287 [sic]; EP 0 389 063). However, these methods are not suitable for obtaining low-endotoxin or endotoxin-free nucleic acid fractions. Thus it has been possible to show, for example, that the measures according to Marko et al. (1982) lead to an endotoxin content of more than 10,000 U per  $\mu\text{g}$  of DNA. Such an endotoxin-rich DNA fraction is unsuitable for transfecting cells in applications of gene therapy.

It was therefore the object of the invention to provide a method for preparing endotoxin-free nucleic acids or nucleic acids with reduced endotoxin content, as a result of which the disadvantages of established methods, such as in particular complicated column materials, are avoided.

The object is achieved by a method for isolating and purifying nucleic acids and/or oligonucleotides from biological samples, in which the particular biological sample is disrupted, undissolved cell components are resuspended in an aqueous potassium acetate solution, optionally present insoluble components are removed, for example by centrifugation, and the aqueous phase is mixed and incubated with an alcoholic solution containing a detergent. The solution is then contacted with a silica gel-like support material, the aqueous

phase is, if possible, quantitatively removed from the support material binding the nucleic acids or oligonucleotides, for example by suction or centrifugation, and the support material with the DNA is then washed adequately. The washing solution used may be an alcoholic solution or acetone which has proved particularly advantageous. Depending on the volume of the starting sample, an incubation time for contacting the support material of from 10 to not more than 40 minutes at room temperature is sufficient; according to the invention, approx. 20 minutes are normally sufficient.

The skilled worker in principle knows silica gel-like support materials. According to the invention, a suspension of silicon dioxide has proved particularly suitable. A silicon oxide suspension which was prepared by adding acid (e.g. hydrochloric acid) to an aqueous suspension of silicon dioxide and was then autoclaved is particularly suitable for the method of the invention.

The aqueous potassium acetate solution contains potassium acetate preferably in a concentration range from approx. 1 to 6 mol/l, and a range from 2 to 4 mol/l and a weakly acidic pH (approx. pH 4.5-6.8) have resulted, according to the invention, in a particularly high quality of the nucleic acids.

Another advantageous embodiment of the method of the invention is to add to the sample, after addition of the potassium acetate solution, additionally one or more RNA-digesting enzymes such as, for example, RNase A and/or RNase T1. In particular for relatively large preparations it has proved advantageous to add the RNA-digesting enzyme(s) in the same medium/buffer in which the potassium acetate salt had been added before. Alternatively, and this is particularly true for relatively small mixtures, the RNA-digesting enzymes

can also be added even during disruption of the biological sample, i.e. together with the lysis buffer (e.g. together with buffer P1 in example 1.2). If a plurality of RNA-digesting enzymes is added, said enzymes may be present in any ratios or else in equal parts. The final concentration of RNA-digesting enzymes in said solution is normally up to or at approx. 150 µg/ml; but even higher enzyme concentrations have not had an adverse effect on the method of the invention.

Normally, according to the invention, an incubation with the potassium acetate solution of from 5 to 10 minutes at 4°C, where appropriate initially at room temperature, is already sufficient for the enzymatic digestion; depending on the amount of sample material used, however, the incubation may also be extended accordingly.

Suitable alcoholic solutions according to the invention are in particular high percentage solutions of higher alcohols such as isopropanol. According to the invention, it has proved particularly advantageous if the alcoholic solution is not diluted with water, that is to say virtually 100% of it consists of the particular alcohol, and it additionally contains one or more ionic detergents, at a concentration of 0.5 to 10% (w/v). A 100% isopropanol solution containing approx. 1 to 4% (w/v) SDS has proved particularly suitable according to the invention.

The biological sample can in principle be disrupted or pre-purified according to methods known to the skilled worker. According to the invention, preference is given to alkaline lysis measures, in particular in the case of bacterial host cells. In this way it is possible to remove protein components and other soluble components before contacting the residue which essentially contains nucleic acid components and other non-soluble

cell components with the potassium acetate solution or the alcohol/detergent solution.

5 Using the method of the invention it is possible to obtain nucleic acids such as, for example, plasmid DNA in high quality, i.e. in particular with an endotoxin content of less than 100 U/ $\mu$ g of DNA, normally of not more than 10 U/ $\mu$ g of DNA.

10 In particular it must be regarded as surprising that the DNA can be bound with high efficiency to the adsorption matrix after alkaline lysis without the need for the addition of chaotropic substances as described in the prior art. The absence of added chaotropic  
15 substances leads to substantial improvements and purifications in the subsequent DNA purification procedure and/or in the corresponding transfection of target cells, that is for cells of both eukaryotic and prokaryotic origin.

20 Moreover, the endotoxin-free nucleic acids and/or oligonucleotides or the nucleic acids and/or oligonucleotides with reduced endotoxin content, which are obtainable according to the method of the  
25 invention, are suitable for producing agents for the treatment of genetic diseases.

The invention further relates to means or compositions for obtaining plasmid DNA from appropriate host cells,  
30 which can be, for example, microtiter plates or blocks which may, where appropriate, contain mini columns for purifying plasmid DNA.

The compositions of the invention essentially contain  
35 an aqueous potassium acetate solution and also a detergent-containing alcoholic solution and a silica gel-like support material. Moreover, it is advantageous if a solution suitable for disrupting a biological sample, in particular for alkaline lysis, is present.



In preferred embodiments of the composition the salt concentration in the potassium acetate solution is in a range from approx. 1 to 6 M, particularly preferably from approx. 2 to 4 M in a weakly acidic medium (pH approx. 4.5-6.8), the alcoholic solution contains isopropanol with approx. 0.5 to 10% (w/v) of an ionic detergent such as, for example, SDS and/or the support material is an aqueous suspension of silicon dioxide.

10 Figure 1

Endotoxin (lipopolysaccharide, LPS) content in various DNA plasmid fractions after acetone washing ((c),(d)) and SDS precipitation ((b),(d)). The plasmid DNA was isolated by binding to silicon oxide and subsequently washed with isopropanol ((a),(b)) or acetone ((c),(d)), with or without LPS precipitation in the presence of SDS (2.5% in isopropanol). The LPS content was determined colorimetrically, according to the manufacturer's instructions (Boehringer Ingelheim, Germany).

- (a) isopropanol/without SDS,
- (b) isopropanol/with SDS,
- (c) acetone/without SDS,
- 25 (d) acetone/with SDS

The following examples further illustrate the invention:

30 1.1 Cell culture and transfection

Baby hamster kidney (BHK) cells were cultured in DMEM (Dulbecco's Modified Eagle's Medium) supplemented with 5% fetal calf serum (Sigma, Deisenhofen, Germany) in a humidified 5% CO<sub>2</sub> atmosphere. For transfections, the cells were applied to 24-well plates and transfected with 2 µg of plasmid DNA according to the calcium phosphate coprecipitation method as described by Roussel et al. (Mol. Cell. Biol. 4 (1984), 1999-2009).

For this purpose, 25 µl of DNA solution were mixed with 25 µl 2 x HBS: 274 mM NaCl, 10 mM KCl, 40 mM HEPES, 1.4 mM Na<sub>2</sub>PO<sub>4</sub>, pH 6.9 at 4°C in a 96-well plate using a 12-channel pipette (Eppendorf, Hamburg, Germany). After adding 20 µl of a 0.25 M CaCl<sub>2</sub> solution (4°C) and mixing, 38 µl were added to the cells after incubation at room temperature for 25 min.

Appropriate aliquots were inoculated in 900 µl of TB medium in wells of 96-well blocks (Qiagen, Hilden, Germany) and cultured with shaking at 300 rpm for approx. 30 hours (37°C). After identification of a positive pool, the DNA was again transfected to confirm the result. The remaining DNA was used to transform bacteria for large-scale plasmid isolation.

#### 1.2 Plasmid isolation with columns

96-well blocks (Qiagen, Hilden, Germany) with bacteria were centrifuged at 3000 g (Sigma centrifuges, Osterode am Harz, Germany) for 5 min. The supernatant was decanted and the blocks were inverted and put on absorbent paper towel for 2 to 3 min. Then 170 µl of buffer P1 (50 mM Tris-HCl/10 mM EDTA pH 8.0, 4°C) were added and the bacteria pellets were resuspended by complete vortex treatment for 10 to 20 min. After addition of 170 µl of buffer P2 (200 mM NaOH, 1% SDS), the block was sealed with foil, inverted and incubated at room temperature for 5 min. The lysis was stopped by adding 170 µl of 4°C cold buffer P3 (3 M potassium acetate pH 5.5, 4°C). Then 10 µl of RnaseA solution (1.7 mg/ml) were added, followed by incubation at room temperature and then at -20°C for 5 min and another centrifugation at 6000 rpm for 10 min. The supernatant was decanted into new blocks and 100 µl of buffer P4 (2.5% (w/v) SDS in isopropanol) were added. The block was subjected to vortexing for 5 min and incubated initially at 4°C for 15 min and then at 20°C for 15 min. The blocks were centrifuged at 6000 rpm for 10 min and

the supernatant was [lacuna] into an array of 96 columns (Qiagen) in appropriately cut 96-well plates, had been prepared [sic]. These plates were placed in vacuum chambers (Qiagen). Then 150 µl of silicon oxide suspension were added followed by incubation at room temperature for 20 min (the silicon oxide suspension was prepared by adding 150 µl of HCl (37%) to 250 ml of a suspension of 50 mg/ml SiO<sub>2</sub> (Sigma) and subsequent autoclaving).

10

After applying reduced pressure, the columns were washed twice with 600 µl of acetone (-20°C). The 96-well column plate was put on a 96-well microtiter plate and centrifuged at 6000 rpm for 4 min. The column plate was dried initially at 37°C for 5 min and then in a vacuum chamber for 5 min and then put on another microtiter plate. 70 µl of double-distilled H<sub>2</sub>O (60°C) were added followed by centrifugation at 6000 rpm for 3 min. The microtiter plate was stored at -20°C.

20

### 1.3 Plasmid isolation without columns

Up to the addition of buffer P4, the method was carried out as described under point 1.2. After centrifugation at 6000 rpm for 10 min, the supernatant was then provided to 96-well POM-microtiter blocks (POM= polyoxymethylene) and 150 µl of silicon oxide suspension were added followed by incubation at room temperature for 20 min. The plates were centrifuged at 6000 rpm for 5 min. The supernatant was carefully decanted and 400 µl of acetone (-20°C) were added. The plates were again vortexed (30 sec) and centrifuged at 6000 rpm for 3 min. This acetone washing was repeated once. The plates were dried initially at room temperature for 5 min and then in a vacuum chamber for 5 min. The pellets were resuspended in 75 µl of water (60°C) and centrifuged at 6000 rpm and 4°C for 10 min. The supernatant was stored in a 96-well microtiter plate at -20°C.

## 2. Results

Plasmid DNA was isolated from the bacteria cultures using mini columns (see point 1.2). A corresponding  
5 protocol without columns is described under point 1.3.

It is important for the transfection step to obtain plasmid DNA of very high purity. For this purpose, silicon dioxide was used as binding matrix for plasmid  
10 DNA. Binding of DNA and silicon dioxide in the presence of chaotropic substances is well known (Vogelstein and Gillespie, Proc. Natl. Acad. Sci. USA 76 (1979), 615-619). Surprisingly, however, it was found that even in the absence of an added chaotropic substance such as,  
15 for example, guanidine hydrochloride, the plasmid DNA binds to silicon dioxide with sufficient capacity. After subsequent washing in acetone, where appropriate with the addition of SDS, plasmid DNA in excellent quality, corresponding to a purification via a cesium  
20 chloride gradient, could be obtained. Commonly, about 10 µg of plasmid DNA with an OD<sub>260/280</sub> of greater than 1.8 were obtained from 900 µl of LB medium, 90% of which were present in supercoiled form.

## 25 3. Comparison with prior art

Experiment A: Bacteria culture: *E.coli* HB101  
pCMVbetaSportGAL, OD<sub>680</sub>/ml approx. 3.3

30 In duplicate mixtures, 1.8 ml each of bacteria culture were worked up using the High Pure plasmid isolation kit (Boehringer Mannheim, Cat. No. 1 754 777), which contains a glass-like support material and a strongly chaotropic salt and 1.8 ml each of bacteria culture  
35 were processed according to the method of the invention.

The result is as follows:

Yield OD <sub>260nm</sub> :	Endotoxin content (LAL assay)
High Pure 1: 9.0 µg/100 µl of endotoxin-free water	214 EU/µg of plasmid
High Pure 2: 8.6 µg/100 µl of endotoxin-free water	240 EU/µg of plasmid
Invention 1: 11.00 µg/100 µl of endotoxin-free water	1.41 EU/µg of plasmid
Invention 2: 10.35 µg/100 µl of endotoxin-free water	4.65 EU/µg of plasmid

Procedure according to the method of the invention using a High Pure filter tube:

- 5 The bacteria culture was centrifuged at 13,000 rpm for 30 sec and the supernatant was removed.

The cell pellet of 1.8 ml of bacteria culture was further treated as follows:

10

1. Resuspending in 250 µl of 50 mM Tris-HCl/10 mM EDTA, 100 µg of RNase (DNase-free), pH 8.0, 4°C.

15

2. Adding 250 µl of 0.2 M NaOH, 1% SDS and 5-10 x inverting the vessel, 5 min at room temperature.

20

3. Adding 250 µl of 3 M K acetate pH 5.5 (4°C) and 5-10 x inverting the vessel, incubating on ice for 5 min.

25

4. Centrifuging in a bench-top centrifuge at maximum speed for 10 min (14,000 rpm), removing the supernatant and adding 0.2 vol. (approx. 150 µl) of 2.5% SDS in isopropanol (e.g. 7 ml of isopropanol and 1 ml of 20% SDS) and vortexing briefly, incubating at 4°C for 15 min and then incubating at -20°C for 15 min.

5. Centrifuging in a bench-top centrifuge at maximum speed for 10 min (14,000 rpm), removing supernatant.
- 5 6. Pipetting supernatant into High Pure filter tube and incubating at room temperature for 20 min.
7. Centrifuging in a bench-top centrifuge at maximum speed for 30 sec (14,000 rpm), discarding the flow-through and washing the filter tube 2 x with 700 µl of ice-cold acetone (centrifuging between the washing steps at 14,000 rpm for 30 sec).
- 10
8. After the last washing step, centrifuging again at 14,000 rpm for 30 sec in order to dry the fleece.
- 15
9. Eluting DNA by adding 100 µl of endotoxin-free water and incubating at room temperature for 10 min. The DNA is obtained by centrifuging at maximum centrifugation speed for 30-60 sec.
- 20

Experiment B: Bacteria culture: *E.coli*  
 JM109pCMVbetaSportGal OD<sub>580</sub>/ml 2.37

Sample	Method	Modification	Yield [µg/ 100µg]	Endotoxin [EU/µg]
1 and 2	High Pure		9.3/9.3	371.7
3 and 4	High Pure	Incubated on fleece for 20 min; incubated before elution for 10 min	12.8/12.2	2.18
5 and 6	Invention	Without incubations	12.2/12.6	0.63

Result:

- The method of the invention shows approx. 100 fold reduction in endotoxin.
- 5 - Furthermore, the inventive method with rapid passing through by centrifugation gives the same yield as using incubation on fleece, and therefore the purification time can now be stated as approx. 70 min. In addition, the inventive method with  
10 rapid passing through by centrifugation shows a lower endotoxin value than after incubation on fleece.

**We claim:**

1. A method for isolating and purifying nucleic acids  
5 and/or oligonucleotides from a biological sample,  
characterized in that
  - the biological sample is disrupted, protein  
10 components and other insoluble components are  
removed,
  - an aqueous solution of potassium acetate is  
15 added to the residue and non-soluble components  
are removed,
  - the potassium acetate-containing solution is  
mixed and incubated with an alcoholic solution  
containing a detergent,
  - the supernatant obtained is contacted and  
20 incubated with a silica gel-like support  
material, and
  - the purified nucleic acids and/or  
25 oligonucleotides are isolated from the soluble  
fraction.
2. The method as claimed in claim 1, characterized in  
30 that the alcoholic solution is a mixture of  
isopropanol with an ionic detergent.
3. The method as claimed in claim 1 or 2,  
35 characterized in that the alcoholic solution  
contains one or more ionic detergents at a  
concentration of 0.5 to 10% (w/v) in 100% strength  
alcohol.



4. The method as claimed in any of claims 1 to 3, characterized in that an aqueous solution containing 1 to 6 M potassium acetate is used.
- 5 5. The method as claimed in claim 4, characterized in that the solution contains 2 to 4 M potassium acetate.
- 10 6. The method as claimed in any of claims 1 to 5, characterized in that the silica gel-like support material used is a suspension of silicon dioxide.
- 15 7. The method as claimed in any of claims 1 to 6, characterized in that the silica gel-like support material is washed with acetone.
- 20 8. The method as claimed in any of claims 1 to 7, characterized in that plasmid DNA with an endotoxin content of less than 100 U/ $\mu$ g is obtained.
- 25 9. The method as claimed in claim 8, characterized in that the endotoxin content is not more than 10 U/ $\mu$ g of plasmid DNA.
- 30 10. An endotoxin-free nucleic acid or oligonucleotide or a nucleic acid or oligonucleotide with reduced endotoxin content obtainable according to a method as claimed in any of claims 1 to 9.
- 35 11. The use of nucleic acids and/or oligonucleotides obtained according to any of the methods as claimed in claims 1 to 9 for transfecting eukaryotic or prokaryotic cells.
12. The use of a nucleic acid and/or oligonucleotides obtained according to any of the methods as claimed in claims 1 to 9 for producing an agent for the treatment of genetic disorders.

13. A composition comprising the following components:
- at least one solution suitable for the disruption of a biological sample,
  - an aqueous potassium acetate solution,
  - a solution of detergent/alcohol, and
  - a silica gel-like support material.
14. The composition as claimed in claim 13, characterized in that the following components are included:
- a solution suitable for alkaline lysis of biological sample material,
  - a salt solution containing 1 to 6 M potassium acetate,
  - an alcoholic solution containing 0.5 to 10% (w/v) SDS in 100% strength isopropanol and
  - a silica gel-like support material.
15. The composition as claimed in claim 13 or 14, characterized in that the support material included is a suspension of silicon dioxide.
16. The use of potassium acetate for isolating, purifying and/or separating endotoxin-free nucleic acids and/or oligonucleotides or nucleic acids and/or oligonucleotides with reduced endotoxin content from and of, respectively, a pre-purified biological sample.

**Abstract**

Method for isolating and purifying nucleic acids and/or oligonucleotides from a biological sample, the use of the isolated or purified nucleic acid and/or oligonucleotide for transfecting cells and also for the production of an agent for the treatment of genetic disorders, a composition suitable for the isolation or purification method and also the use of potassium acetate and a silica gel-like support material for isolating endotoxin-free nucleic acids and/or oligonucleotides or nucleic acids and/or oligonucleotides with reduced endotoxin content.